

EPC/GDV MEASUREMENTS DURING SUN ECLIPS 01.08.2008 IN SIBERIA

1. SUN ECLIPSE INFO

Source: <http://www.eclipse-2008.ru/eclipse.php>

Sun Eclipse 1 August 2008 has N 126 in accordance with clasification. First Sun Eclipse of this series was in 1179, but only from 1882 a series of full sun eclipses began 1882, 1900, 1918, 1936, 1954, 1972, 1990, 2008, 2026, 2044. Series ends in 2459.

The Map of Sun Eclipse 01 August 2008




Table 1 shows parameters for several sites. Time difference from Grinvich time is 7 hours.

P1 - the moment of the first contact; U1 - the moment of the second contact (beginning of a full phase); U2 - the moment of the third contact (the end of a full phase); P2 – the moment of the fourth contact

Site	Long., min	Max	Phase	P1	U1	U2	P2
Novosibirsk +55 02/82 55	2:21	17 45 9	1,017	16 41 17	17 43 58	17 46 20	18 45 7
Ob +55 0/82 43	2:22,9	17 45 9	1,019	16 41 13	17 43 58	17 46 21	18 45 9
Berdsk +54 47/83 2	2:22,1	17 45 40	1,018	16 41 49	17 44 29	17 46 51	18 45 35

2. PROTOCOL OF MEASUREMENTS

Equipment:

EPC/GDV Camera with battery power supply.

Computer with any type of power supply.

Sensor – antenna.

Air fan – home appliance or computer.

Protocol of measurements:

1. Metal cylinder – test-object is positioned on the electrode. The wire is fixed with a tape – any movement of the test-object will distort results. Fan is placed nearby the Camera for constant ventilation.
2. Instrument warming up. Test-object is connected with grounding jack of the Camera. Regime 1, series of images, 3 s interval, during 1 hour.
3. Test-object wire is connected with Antenna.
4. Background measurement. Regime 1, series of images, 3 s interval, during 1 hour.
5. Measurement during and after solar eclipse in the same regime during 1 hour or more.

Processing: “GDV SciLab” program.

3. ORGANISATION OF THE EXPERIMENT

In the process 6 EPC/GDV Cameras have been used.

Positions:

1. Novosibirsk Academic City – Hypo-magnet Camera.
2. Novosibirsk Academic City – placebo of the Hypo-magnet Camera (camera without permalloy layer).
3. Novosibirsk – countryside.
4. Berdsk
5. Irkutsk
6. Novosibirsk city center.

Overall organization: K. Korotkov and Oleg Sorokin.

Researchers:

Research in the Novosibirsk Academic City was done in the frame of the big program under the leadership and with the help of professor Alexander Trofimov.

The protocol of measurements in Novosibirsk city center was different from all other sites and results are not included in this report.

4. RESULTS

Measurements with antenna started at about 16.30. Signal was very stable for all the measured sites: variability of data was about 3% before the sun eclipse and less than 1% after the sun eclipse. Significant variations in the precise structure of a signal was recorded.

Graphs 1-7 demonstrate time dynamics of Area changes, graphs 8-14 time dynamics of Intensity. On the abscise axis the N of point and absolute time in Novosibirsk is given: 163004 = 16 h 30 min 04 sec. Arrows roughly mark the time of full sun eclipse (1743 on Novosibirsk time).

As we see from the graphs, they all have two clear phases: before and after the sun eclipse. Before sun eclipse all graphs demonstrate longitudinal waves. After the sun eclipse signal stabilizes. Two types of dynamics before the sun eclipse are recorded: decreasing (for two instruments) and increasing (for five instruments).

Time dynamics of graphs fig 1 and 2 are quite similar. There are correlations of position of three local extremum. At about 17.39 an increased variability of data is observed.

Graphs of fig. 3-5 do not have decreasing trend typical for graphs fig. 1 and 2, but they have local extremums as well. Rough position of the extremis are given in Table 2. As we see from Table 2, the position of some extremums are quite similar.

Table 2. Rough position of the extremis on the Area graphs before the sun eclipse.

Place	Hypo-magnet	Placebo	Countryside	City Center	Berdsk	Abakan
Rough Time of max		16_30		16-21	16-31	16-32
	16-35	16_35	16-37		16-37	16-38
	16-45	16_46	16-42	16-50		16-50
		16-55			17_02	17_02
	17-15					
	17-25			17-23		17-19

GRAPHS OF AREA DYNAMICS




Fig.1. Area time dynamics in the Placebo of the Hypo-magnet Camera




Fig.2. Area time dynamics in Berdsk




Fig. 3. Area time dynamics in Hypo-magnet Camera




Fig.4. Area time dynamics in the countryside




Fig. 5. Area time dynamics in Irkutsk

In Abakan the amplitude of a signal changed abruptly at 17.35. We do not know the reason of this situation so we present graphs before and after the sun eclipse separately.




Fig.6. Area time dynamics in Abakan before the sun eclipse.




Fig.6. Area time dynamics in Abakan after the sun eclipse.

Time dynamics of the Intensity




Fig. 8. Time dynamics of Signal Intensity in Hypo-magnet Camera




Fig. 9. Time dynamics of Signal Intensity in Placebo Hypo-magnet camera




Fig.10. Time dynamics of Signal Intensity in Berdsk




Fig.11. Time dynamics of Signal Intensity in countryside




Fig.12. Time dynamics of. Signal intensity in Irkutsk




Fig.13. Time dynamics of. Signal intensity in Abakan before the sun eclipse




Fig.14. Time dynamics of. Signal intensity in Abakan after the sun eclipse

Fig. 15 and Table 3 demonstrate data on Area and Intensity variability before and after the sun eclipse (ratio between Standard Deviation and Average in %).

Table 3. Data Variability

	Before	After	Before	After	Before	After	Before	After	Before	After	Before	After
Area	Hypo-magnet		Placebo		Countryside		Berdsk		Irkutsk		Abakan	
Average	3778.9	3829.1	7719.5	7411.2	6929.1	6988.7	4482.1	4302.0	8325.7	8632.3	12933.2	3443.2
St. Deviation	111.4	51.55	275.22	104.39	141.21	141.16	302.46	73.19	228.7	89.9	234.4	55.8
Variability	0.029	0.013	0.036	0.014	0.020	0.020	0.067	0.017	0.0275	0.0104	0.0181	0.0162
Intensity	Hypo-magnet		Placebo		Countryside		Berdsk		Irkutsk		Abakan	
Average	57.954	57.84	57.22	58.06	91.20	93.33	59.23	60.25	96.7	100.4	113.7	56.7
St.Deviation	1.347	0.44	1.01	0.74	2.41	0.83	1.45	0.48	2.3	0.6	3.9	0.4
Variability	0.023	0.008	0.018	0.013	0.026	0.009	0.025	0.008	0.0239	0.0062	0.034	0.008





Fig.15. Area and Intensity data variability before (blue bars) and after (red bars) the sun eclipse.

CONCLUSIONS

1. Seven EPC/GDV positioned in different locations recorded statistically different signal in different phases of sun eclipse. Statistically significant difference of readings before and after with probability 99.99999 ($p < 0.00001$) both for Area and Intensity was found. For other parameters the difference was clear visible, but was not statistically significant in all cases.
2. Before sun eclipse longitudinal waves was recorded, their origin should be found.
3. Position of extremums concur for several EPC instruments. Some extremums are different.
4. Two instruments demonstrated signal with decreasing amplitude before the sun eclipse, other instruments demonstrated increasing trend of a signal before sun eclipse.
5. The highest amplitude of oscillations was found for the instrument positioned in the hypo-magnet camera.
6. In the phase after the sun eclipse all the instruments recorded very stable signal with variability less than 1%.
7. It is recommended to make similar measurements for longer time before the sun eclipse.
8. Presented results open up perspectives for EPC/GDV instruments applications for early registration of geophysical anomalies.

We appreciate your comments.

Thank you

Dr. Konstantin Korotkov kk@korotkov.org

References

- Korotkov K., Korotkin D. Concentration dependence of gas discharge around drops of inorganic electrolytes. J of Applied Physics, 2001, 89, 9, 4732-4737.
- Measuring Energy Fields: State of the Art. GDV Bioelectrography series. Vol. I. Korotkov K. (Ed.). Backbone Publishing Co. Fair Lawn, USA, 2004.
- Korotkov K., Krizhanovsky E., Borisova M., Hayes M., Matravers P., Momoh K.S., Peterson P., Shiozawa K., and Vainshelboim A. The Research of the Time Dynamics of the Gas Discharge Around Drops of Liquids. J of Applied Physics. 2004, v. 95, N 7, pp. 3334-3338.
- Bell I., Lewis D.A., Brooks A.J., Lewis S.E., Schwartz G.E. Gas Discharge Visualisation Evaluation of Ultramolecular Doses of Homeopathic Medicines Under Blinded, Controlled Conditions. J of Alternative and Complementary Medicine, 2003, 9, 1: 25-37